

Title	The influence of S-2-aminoethyl isothiuronium Br-HBr on the equal effect ratio between 200kVp X-rays and 60Co $\gamma$ -rays
Author(s)	岡村, 重昭; 馬屋原, 晟; 古賀, 充 他
Citation	日本医学放射線学会雑誌. 27(3) p.253-p.260
Issue Date	1967-06-25
oaire:version	VoR
URL	<a href="https://hdl.handle.net/11094/14789">https://hdl.handle.net/11094/14789</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

# THE INFLUENCE OF S-2-AMINOETHYL ISOTHIURONIUM Br-HBr ON THE EQUAL EFFECT RATIO BETWEEN 200kVp X-RAYS AND $^{60}\text{Co}$ $\gamma$ -RAYS

by

Shigeaki Okamura

Department of Experimental Radiology (Director: Prof. H. Yoshinaga),  
Faculty of Medicine, Kyushu University

Akira Umayahara Mitsuru Koga, Toshikazu Takeshita and Akira Yasukochi  
Department of Radiology (Director: Prof. H. Irie), Faculty of Medicine,  
Kyushu University, Fukuoka, Japan

200 kVp X 線と  $^{60}\text{Co}$   $\gamma$  線の等効果比に及ぼす S-2-Amino-  
ethylisothiuronium Br-HBr の影響

九州大学医学部放射線基礎医学教室（指導 吉永春馬教授）

岡 村 重 昭

九州大学医学部放射線科学教室（指導 入江英雄教授）

馬屋原 晟 古賀 充 安河内 彰 竹下 寿七

（昭和42年10月20日受付）

雄性及び雌性の CF#1 10~11週令のハツカネズミを用い、その  $\text{LD}_{50/30}$  を指標にした 200 kVp X線と  $^{60}\text{Co}$   $\gamma$ 線の等効果比に及ぼす S-2-Aminoethylisothiuronium Br-HBr（以後 A E T と略する）の影響を検討した。

$\text{LD}_{50/30}$  は X線群の雄 560 rads, 雌 553 rads, X線 + A E T 処置群で雄 810 rads, 雌 902 rads,  $\gamma$ 線群で雄 874 rads, 雌 857 rads,  $\gamma$ 線 + A E T 処置群で雄 1248 rads, 雌 1325 rads となった。

これより A E T の dose-reduction factor を求めると X線群で雄 1.5, 雌 1.6,  $\gamma$ 線群で雄 1.4, 雌 1.5 となり、等効果比を求めると A E T 非投与

群では雄 0.6, 雌 0.6 : A E T 投与群では雄 0.6, 雌 0.6 となった。

この結果から考えると A E T の dose-reduction factor は X線でも  $\gamma$ 線でもほぼ等しく、又 A E T の投与は X線と  $\gamma$ 線の等効果比に影響を及ぼさないと考えられる。

何故 A E T が X線と  $\gamma$ 線の等効果比に影響しなかったかについて考察し、通常用いられている Gray や Cormack の計算した L E T 値よりも Burch の計算した L E T 値の方がより適切であろうと述べた。

## Introduction

Numerous reports have been made on the protective agents against radiation injuries. Since the report of Doherty et al.<sup>1,2)</sup> S-2-Aminoethylisothiuronium Br-HBr (AET) in particular has been recognized to be the most effective protective agent against radiation damage.

On the other hand, many experiments have been conducted on the relative biological effectiveness (RBE) between radiations of different qualities.<sup>3)</sup> Particularly, the RBE between 200 kVp X-rays and high energy radiations has been studied by Irie and his coworkers with the support of the scientific research grant provided by the Ministry of Education of Japan<sup>4)5)6)7)</sup>.

However, there is hardly any report in literature which compares the effect of protective agents against radiations of different qualities.

In the present study the difference in effect of AET on the LD<sub>50/30</sub> of mice between 200 kVp X-rays and <sup>60</sup>Co γ-rays was investigated.

### Experimental Procedure

CF #1 male and female mice, 10 to 11 weeks old, were divided into the following four groups. Each exposure was graded in these four groups, as described below.

X-ray groups (for both sexes)	480, 550, 630, and 725 R
X-ray+AET groups (for male)	630, 725, 800, 900, and 1000 R
(for female)	800, 900, 1000, and 1100 R
X-ray groups (for both sexes)	725, 800, 900, and 1000 R
X-ray+AET groups (for male)	1100, 1200, 1300, 1400, and 1500 R
(for female)	1200, 1300, 1400, and 1500 R

Radiation sources and exposure conditions are as follows.

200 kVp X-rays: Shinai X-ray machine (manufactured by Shimazu Co.) operated at 15 mA with a filtration of 0.7 mm Cu+0.5 mm Al. The distances from X-ray target to the mouse were 55 cm and 33 cm and dose rates were 52.5 R/min and 151.0 R/min for each distance, as measured in the air with a Victoreen ionization chamber (No. 154).

<sup>60</sup>Co γ-rays: 1.17 MeV and 1.33 MeV γ-rays from a 3000 -curie teletherapy unit (Shimazu). The distances from the source to the mouse were 106 cm and 67 cm and dose rates were 46.7 R/min and 128.0 R/min for each distance, as measured in the air with a Victoreen chamber (No. 621).

The high dose rates of X-ray and γ-ray were used in the cases irradiated over 800 R or 900 R. Whole body irradiation was delivered in a rectangular wooden container, covered with X-ray film for X-ray irradiation and with acrylate resin plate (thickness of 5 mm) for γ-ray, respectively.

A dose of 7 mg of AET neutralized by 0.1 N NaOH (namely, AET changes to Mercaptoethyl Iguanidine in this condition) was injected intraperitoneally and the corrected administration time<sup>8)</sup> ("time from administration to irradiation" +  $\frac{\text{irradiation time}}{2}$ ) was 20 minutes.

### Results

The dose, the number of mice, the number surviving 30 days after exposure, and the survival rate (probit) are shown in Table I.

Attempts were made to develop an empirical transformation formula which would give values decreasing linearly with irradiation dose. A modified logarithmic transformation formula was found to be satisfactory.

$$S = -aD + b$$

where D is irradiation dose in log. scale, S is the most reliable survival ratio in the probit scale, and a

Fig. I. The relationship between exposure and survival rate (male) is illustrated.





Table I. The doses, the number of mice, the number surviving 30 days after exposure, and the survival rate (probit) are shown.

Male					Female				
X-ray groups									
Dose	No. of mice	30 days survival	survival rate	probit	No. of mice	30 days survival	survival rate	probit	
480R	12	10	83.3%	5.97	32	29	90.6%	6.32	
550R	13	9	69.2%	5.50	33	25	78.8%	5.80	
630R	11	4	36.4%	4.65	33	8	24.2%	4.30	
725R	14	2	14.3%	3.93	29	1	3.4%	3.17	
X-ray + AET groups									
630R	14	11	78.6%	5.79					
725R	13	8	61.5%	5.30					
800R	13	8	61.5%	5.30	33	28	84.8%	6.03	
900R	15	8	53.3%	5.09	36	27	75.0%	5.67	
1000R	15	4	26.7%	4.38	35	11	31.4%	4.51	
1100R					35	5	14.3%	3.93	
gamma-ray groups									
725R	11	10	90.9%	6.33	28	26	92.9%	6.47	
800R	11	9	81.8%	5.91	31	27	87.1%	6.13	
900R	14	6	42.9%	4.82	33	19	57.6%	5.19	
1000R	12	4	33.3%	4.57	33	4	12.1%	3.82	
gamma-ray + AET groups									
1100R	12	10	83.3%	5.96					
1200R	14	8	57.1%	5.18	34	28	82.4%	5.93	
1300R	14	6	42.9%	4.82	34	24	70.6%	5.54	
1400R	14	5	35.7%	4.63	36	13	36.1%	4.64	
1500R	12	4	33.3%	4.57	30	10	33.3%	4.57	

Fig. II. The relationship between exposure and survival rate (female) is illustrated.



and b are constant.

This linearity in relationship between survival ratio and exposure is demonstrated in Figure I and II. The relationship between survival ratio and log. dose is calculated <sup>9)</sup>as follows.

Male

X-ray groups  $S = -11.698D + 37.41$

X-ray + AET groups  $S = -6.111D + 22.91$

γ-ray groups  $S = -13.588D + 45.21$

γ-ray + AET groups  $S = -10.053D + 36.30$

Female

X-ray groups  $S = -18.381D + 55.82$

X-ray + AET groups  $S = -16.215D + 53.01$

γ-ray groups  $S = -18.935D + 60.86$

γ-ray + AET groups  $S = -15.462D + 53.55$

In the above, D is roentgen in log. scale, S is the most reliable value of survival ratio in the probit scale.

Table II LD<sub>50/30</sub> values of CF#1 male and female mice on X-ray groups, X-ray + AET groups, γ-ray groups, and γ-ray + AET groups are shown.

	male	female
X-ray groups	590 R ( 560 rads)	582 R ( 553 rads)
X-ray + AET groups	853 R ( 810 rads)	949 R ( 902 rads)
γ-ray groups	910 R ( 874 rads)	891 R ( 856 rads)
γ-ray + AET groups	1299 R (1247 rads)	1380 R (1325 rads)

Table III The dose-reduction factor on X-ray and on  $\gamma$ -ray is shown.

$\frac{LD_{50/30} \text{ of X-ray + AET groups}}{LD_{50/30} \text{ of X-ray groups}}$	1.4 (male)	1.6 (female)
$\frac{LD_{50/30} \text{ of } \gamma\text{-ray + AET groups}}{LD_{50/30} \text{ of } \gamma\text{-ray groups}}$	1.4 (male)	1.5 (female)

Table IV. The equal effect ratio between X-ray and  $\gamma$ -ray with or without AET treatment is shown.

$\frac{LD_{50/30} \text{ of X-ray groups}}{LD_{50/30} \text{ of } \gamma\text{-ray groups}}$	0.6 <sub>4</sub> (male)	0.6 <sub>5</sub> (female)
$\frac{LD_{50/30} \text{ of X-ray + AET groups}}{LD_{50/30} \text{ of } \gamma\text{-ray + AET groups}}$	0.6 <sub>5</sub> (male)	0.6 <sub>8</sub> (female)

With this formula,  $LD_{50/30}$  of each group was calculated and is shown in Table II.

In this calculation, factors used in converting roentgens to rads were 0.95<sup>10)</sup> for X-ray and 0.96<sup>11)</sup> for  $\gamma$ -ray, respectively.

The dose-reduction factor of AET is shown in Table III.

The dose-reduction factor of AET for X-ray is 1.4 for male and 1.6 for female, and for  $\gamma$ -ray it is 1.4 for male and 1.5 for female, respectively.

The equal effect ratio between X-ray and  $\gamma$ -ray is shown in Table IV.

The equal effect ratio between X-ray and  $\gamma$ -ray without AET treatment is 0.6<sub>4</sub> for male and 0.6<sub>5</sub> for female and with AET treatment it is 0.6<sub>5</sub> for male and 0.6<sub>8</sub> for female, respectively.

### Discussion

The RBE Committee has recommended that the term "equal effect ratio" should be used instead of "RBE" for the comparison of effects between radiations of different qualities, such as  $LD_{50/30}$ . Because  $LD_{50/30}$  is a measure of effects which, although equal in number of fatalities one month after irradiation, can involve very different mechanisms leading to death.

Many authors have made studies to determine the optimum administration time of AET. Some give 30 minutes<sup>13)</sup> before irradiation as the best administration time of AET, while others give 10 minutes<sup>14)</sup> and 10–15 minutes<sup>15)</sup> as the best administration time of MEG, but their irradiation time (or dose rate) differs. The length of irradiation time bears upon the survival rate in the use of protective agents, because the distribution of protective agent in the critical organs changes after administration even during irradiation. Therefore, the length of irradiation may affect the optimum administration time. In 1961 the authors proposed<sup>8)</sup> the corrected administration time which takes consideration into interval time between administration and irradiation (the administration time) and the interval time between the beginning and end of irradiation (the irradiation time). In 1965 this concept of corrected administration time was supported by the report on "the distribution of radioactivity following AET-<sup>35</sup>S administration"<sup>16)</sup>. Thus, the problem of how to determine the administration time in the cases of different irradiation time (or of different irradiation dose) can be resolved by the use of this corrected administration time. In practice, two dose rates were used in order to reduce the difference between the longest and shortest irradiation time.

According to the recent reports made by Oldfield et al.<sup>17)18)</sup> on the chemical protection against 440 MeV protons in mice pretreated with Mercaptoethylamine (MEA) and p-Aminopropiophenone (PAPP),



The indirect action of X-rays and  $\gamma$ -rays is assumed to be chiefly due to  $\delta$ -rays and it is considered that  $\delta$ -rays of both X-rays and  $\gamma$ -rays do not give any qualitative difference in effect, since  $\delta$ -ray has the same character with regard to both radiations. In the mean LET computed by Gray and Cormack  $\delta$ -track formation is ignored. According to Burch<sup>21)</sup> the mean LET values allowing for  $\delta$ -tracks are 25.8 keV/micron for 200 kVp X-rays and 19.6 keV/micron for  $^{60}\text{Co}$   $\gamma$ -rays. Based on our experiment, the mean LET values allowing for  $\delta$ -tracks by Burch are considered to be more reasonable than those calculated by Gray and Cormack.

### Summary

A study was made on the equal effect ratio between 200 kVp X-rays and  $^{60}\text{Co}$   $\gamma$ -rays based on  $\text{LD}_{50/30}$  of CF $\#$ 1 male and female mice with or without S-2-Aminoethylisothiuronium Br-HBr treatment.

The problem of length of irradiation, which influences the survival ratio when protective agents are used, was solved by using the corrected administration time. This was reported earlier.

The dose-reduction factor of AET on X-ray is 1.4 for male and 1.6 for female, and on  $\gamma$ -ray it is 1.4 for male and 1.5 for female, respectively.

The equal effect ratio between 200 kVp X-rays and  $^{60}\text{Co}$   $\gamma$ -rays without AET treatment was 0.6<sub>4</sub> for male and 0.6<sub>5</sub> for female and with AET treatment 0.6<sub>5</sub> for male and 0.6<sub>8</sub> for female.

These results suggested that the dose-reduction factor of AET on X-ray and  $\gamma$ -ray is almost equal and that the equal effect ratio between X-ray and  $\gamma$ -ray is almost equal regardless of AET treatment.

Discussion was made on the reason why AET treatment has not influenced the equal effect ratio and why the mean LET values calculated by Burch are more appropriate than those reported by Gray and Cormack.

### Acknowledgement

This work was supported with the scientific research grant from the Ministry of Education of Japan.

The authors wish to express their sincere thanks to Professor Hideo Irie and Professor Haruma Yoshinaga for their constant guidance and wish to express their appreciation to Assistant-professor Masaharu Kojima (Faculty of Pharmaceutical Sciences, Kyushu University) for the production of AET.

### References

- 1) David G. Doherty, W.T. Burnett, Jr., and Raymond Shapira: Chemical protection against ionizing radiation. II. Mercaptoalkylamines and related compounds with protective activity. *Radiation Research* 7, 13-21, 1957.
- 2) Raymond Shapira, David G. Doherty, and W.T. Burnett, Jr.: Chemical protection against ionizing radiation. III. Mercaptoalkylguanidines and related isothiuronium compounds with protective activity. *Radiation Research* 7, 22-34, 1957.
- 3) John B. Storer, Payne S. Harris, John E. Furchner, and Wright H. Langham: The relative biological effectiveness of various ionizing radiation in mammalian systems. *Radiation Research* 6, 188-288, 1957.
- 4) Hideo Irie et al.: Ann. Rep. Minist. Educ. 1961 Radiation Effects-Coop. Res. p. 77-82 (in Japanese).
- 5) Hideo Irie et al.: Ann. Rep. Minist. Educ. 1962 Radiation Effects-Coop. Res. p. 49-56 (in Japanese).
- 6) Hideo Irie et al.: Ann. Rep. Minist. Educ. 1964 Radiation Effects-Coop. Res. p. 95-107 (in Japanese).
- 7) Hideo Irie et al.: Ann. Rep. Minist. Educ. 1965 Radiation Effects-Coop. Res. p. 81-91 (in Japanese).
- 8) Shigeaki Okamura, Hitoshi Katayama and Seichi Yoshimoto, The advocacy of the "action curve" and the "corrected administration time" of protective agent:—Special experiment of MEG HBr and MEG 1/2  $\text{H}_2\text{SO}_4$ —. *Nipp. Act. Radiol.* 21, 773-779, 1961. (in Japanese).
- 9) George W. Snedecor: STATISTICAL METHODS Applied to Experiments in Agriculture and Biology, p. 104, 1955. Iwanami Press, Tokyo.
- 10) Haruma Yoshinaga, Shigetoshi Antoku, Keiko Tanaka and Yaeko Oku: On the quality of X-ray used in the



- radiological studies. Proc. Hiroshima Univ. RINMB No. 2, p. 11-22, 1961.
- 11) Kenji Takeshita, Seiichi Yoshimoto and Shigetoshi Antoku: Absorbed dose measurements of high energy radiations using ferrous sulfate dosimeter. Nipp. Act. Radiol. 23, 288-292, 1963. (in Japanese).
  - 12) The RBE Committee: Report of the RBE Committee to the International Commissions on Radiological Protection and on Radiological Units and Measurements. Health Physics 9, 357-384, 1963.
  - 13) Ryo Katsuhara: An experimental study of certain new chemicals as protective agents against radiation injury. Nipp. Act. Radiol. 19, 73-92, 1959. (in Japanese).
  - 14) Nobuyoshi Muta and Seichirō Koiwai: The Protective effect of MEG against X-irradiation. Nipp. Act. Radiol. 23, 346-353, 1960. (in Japanese).
  - 15) Shigetoshi Antoku: Chemical protection of MEG against ionizing radiation. Nipp. Act. Radiol. 23, 666-673, 1963. (in Japanese).
  - 16) Hitoshi Katayama: Action of Aminoethylisothiuronium HBr (AET)—Distribution of radioactivity following AET-<sup>35</sup>S administration and optimum administration time of AET. Nipp. Act. Radiol. 25, 124-136, 1965. (in Japanese).
  - 17) D.G. Oldfield, J. Doull, and V. Plzak: Chemical protection against 440 MeV protons in mice pretreated with Mercaptoethylamine (MEA) or p-Aminopropiophenone (PAPP). Radiation Research 26, 12-24, 1965.
  - 18) D.G. Oldfield, J. Doull, and V. Plzak: Chemical protection against absorber-moderated protons. Radiation Research 26, 25-31, 1965.
  - 19) L.H. Gray: The distribution of the ions resulting from the irradiation of living cells. Brit. J. Radiol. Suppl. 1, p. 7-15, 1947.
  - 20) D.V. Cormack and H.E. Johns: Electron energies and ion densities in water irradiated with 200 KeV, 1 MeV and 25 MeV radiation. Brit. J. Radiol. 25, 369-381, 1952.
  - 21) P.R.J. Burch: Some physical aspects of relative biological efficiency. Brit. J. Radiol. 30, 524-529, 1957.
-